

Perturbation study of the conductance through a finite Hubbard chain

Akira Oguri¹

Department of Material Science, Faculty of Science, Osaka City University, Osaka 558-8585, Japan

Abstract

The dc conductance through a Hubbard chain of size N ($=1, 2, 3, \dots$) connected to noninteracting leads is studied at $T = 0$ in an electron-hole symmetric case using a perturbation theory in U . The result shows a typical even-odd property corresponding to a Kondo or Mott-Hubbard physics.

Keywords: quantum transport; electron correlation; Fermi liquid; mesoscopic system

Motivated by a current interest in effects of electron correlation on the transport through small systems, we have examined some theoretical approaches [1,2]. In this report, using a perturbation approach, we study the size (N) dependence of the transport through a small interacting chain connected to semi-infinite leads.

We start with the Hamiltonian $\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_I$,

$$\mathcal{H}_0 = - \sum_{i=-\infty}^{+\infty} \sum_{\sigma} t_i \left(c_{i+1\sigma}^{\dagger} c_{i\sigma} + c_{i\sigma}^{\dagger} c_{i+1\sigma} \right) - \mu \sum_{i=-\infty}^{+\infty} \sum_{\sigma} n_{i\sigma} + \sum_{j=1}^N \left(\epsilon_0 + \frac{U}{2} \right) n_{j\sigma}, \quad (1)$$

$$\mathcal{H}_I = U \sum_{j=1}^N \left[n_{j\uparrow} n_{j\downarrow} - \frac{1}{2} (n_{j\uparrow} + n_{j\downarrow}) \right]. \quad (2)$$

Here $c_{i\sigma}^{\dagger}$ is a creation operator for an electron with spin σ at site i , $n_{i\sigma} = c_{i\sigma}^{\dagger} c_{i\sigma}$. The hopping matrix element is uniform $t_i = t$ except at the boundaries between the central region and two leads; $t_0 = v_L$ and $t_N = v_R$.

At $T = 0$, the dc conductance g_N can be written in term of an inter-boundary element of a single-particle Green's function $G_{N1}(\omega + i0^+)$, and is determined by the value at the Fermi level $\omega = 0$ as $g_N = (2e^2/h) 4 \Gamma_L(0) \Gamma_R(0) |G_{N1}(i0^+)|^2$ [1]. Here $\Gamma_{\alpha}(0) = \pi D(0) v_{\alpha}^2$ with $\alpha = L, R$, and $D(0) = \sqrt{4t^2 - \mu^2} / (2\pi t^2)$. The assumption made here is the validity of the perturbation theory in U . This seems to be probable for small N . In that case, the self-energy due to $\mathcal{H}_I$ has a property $\text{Im} \Sigma_{jj'}(i0^+) = 0$ at $T = 0$ [3], and g_N can be obtained through a scattering problem of free quasi-particles [2].

In this report, we consider an electron-hole symmetric case taking the parameters to be $\mu = 0$ and $\epsilon_0 + U/2 = 0$. If the system has an additional in-

¹ E-mail: oguri@sci.osaka-cu.ac.jp




Fig. 1. g_N vs M for even N ($= 2M$): $U/(2\pi t) = 1.0$, and $v_L/t = v_R/t = 0.7$ ($\star$), 0.8 ($\blacklozenge$), 0.9 ($\blacktriangle$), 1.0 ($\blacksquare$).

version symmetry $v_L = v_R$, it can be shown that a perfect transmission occurs $g_N \equiv 2e^2/h$ for odd N ($= 2M + 1$) independent of the values of U and M [2]. This is caused by the Kondo resonance appearing at the Fermi level for odd N .

On the other hand, for even N , we evaluate the self-energy $\Sigma_{jj'}(i0^+)$ within the second order in U at $T = 0$, and then obtain $G_{N1}(i0^+)$ solving the Dyson equation in the real space [2]. In Fig. 1, g_N for even N ($= 2M$) is plotted vs M for several values of v_L ($= v_R$), where $U/(2\pi t) = 1.0$. The dc conductance decreases with increasing the size $2M$. This behavior can be regarded as a tendency toward a Mott-Hubbard insulator, and it is pronounced for larger U . In Fig. 2, g_N for even N ($= 2, 4, \dots$) is plotted vs U . The value of g_{2M} decreases with increasing U . When v_L (or v_R) is smaller than t , the reduction of g_{2M} is proportional to U^2 at $U/(2\pi t) \ll 1$. As it is seen in the plots for $v_L = v_R = 0.7t$ (dashed lines), the peak structure in the U dependence becomes sharp for large M , and in the limit $M \rightarrow \infty$ the peak seems to vanish leaving the value at a singular point $U = 0$ unchanged. In contrast, in the case of $v_L = v_R = t$ (solid lines), the reduction of g_{2M} is proportional to U^4 at $U/(2\pi t) \ll 1$, and g_{2M} seems to be finite in the limit of large M . However, in order to verify this behavior for large M , the contributions of the higher-order terms should be examined because the unperturbed Hamiltonian $\mathcal{H}_0$ has a translational invariance accidentally in this case.




Fig. 2. g_N vs U for even N ($= 2, 4, \dots$): $v_L/t = v_R/t = 0.7$ (dashed lines), 1.0 (solid lines). 15+30 lines are plotted.

Qualitatively, the even-odd property seems to be understood from that in the unperturbed system, especially from the level structure of the isolated chain. For odd N , there is a semi-occupied one-particle state at the Fermi level $\omega = 0$, and thus a doublet ground state is realized. When the leads are connected, the ground state is replaced by a Kondo singlet state and contributes to the tunneling. On the other hand, for even N , the Fermi level lies between the highest occupied level and the lowest unoccupied level, and thus a finite energy corresponding to the gap is necessary to excite the electrons. Although the levels are broadened by the coupling with the leads, the even-odd property is determined by whether there exists a zero-energy excitation or not. At finite temperatures, a characteristic energy scale of the Kondo or Mott-Hubbard physics will play an important role.

We expect that the model and approach used here can be applied to a series of quantum dots or a quantum wire of nanometer scale.

We thank H. Ishii for valuable discussions.

References

- [1] A. Oguri, Phys. Rev. B **56**, 13422 (1997); **58** 1690(E) (1998).
- [2] A. Oguri, Phys. Rev. B **59**, 12240 (1999); unpublished.
- [3] J. S. Langer and V. Ambegaokar, Phys. Rev. **121**, 1090 (1961).